



REDEFINING PERSONALIZATION IN FIXED BEARING PARTIAL KNEE DESIGN

Zimmer Biomet is the leading company in partial knee arthroplasty¹⁻³ with over 40 years of experience, offering a comprehensive range of anatomic and innovative PKA solutions.

Persona Partial Knee offers:

- Personalized, compartment-specific implant shapes based on the Persona® Total Knee
- Precise, efficient instrumentation
- Proven Vivacit-E Vitamin-E[®] Polyethylene technology⁴⁻⁸

Clinically Proven Legacy^{9,10}

The Persona Partial Knee carries forward design elements of the Zimmer Miller Galante (M/G) Uni, which showed 98% survivorship at 10 years and 90% and at 20 years.⁹

643 Persona Partial Knees were implanted, and two-year results demonstrated:¹¹

- 98.9% survivorship at two years
- 97.3% of patients were satisfied or very satisfied with the results of surgery


PRECISE. PERSONALIZED. PROVENI.

Femoral Component

- Compartment-specific shape
- Consistent AP growth and shape throughout seven femoral sizing options
- Extended posterior condyle safely accommodates high flexion up to 155°



Yellow outline indicates the Zimmer M/G Uni



Tibial Component

- Compartment-specific shape with seven anatomic tibial profiles
- Two peg and keel fixation that has a proven^{9,10} track record going back to the Zimmer M/G Uni*

Persona Partial Knee surgeon developers and engineers studied the morphology of thousands of bones, representing a diverse global population, using ZiBRA™ virtual resection technology. Using this technology led to:

- Improved sizing options and shape of the tibia aim for a better fit, with decreased overhang medially and decreased underhang posteriorly, compared to previous designs¹²
- The anatomic shape of the tibial implant increases tibial coverage¹² aimed at preventing subsidence



PRECISE, EFFICIENT INSTRUMENTATION

Precise and efficient instrumentation facilitates accurate alignment and under-correction of the leg with a simple spacer block technique.



Tibial Resection Guide

- The tibial cut guide allows the surgeon to place the guide on the tibia and orient the vertical saw blade in the preferred M/L location and rotation of the cut
- Tibial resection guides attach to the Persona EM tibial resection tube body



Femoral Finishing Guide

- Persona Partial PK Handle eases insertion
- Extended saw slots allow controlled posterior and chamfer resections
- Three fixation screws placed anteriorly on the guide add stability, allowing resections to be made without removing screws



Locking Femoral Impactor

 Femoral insertion notches were designed for precise control of femoral component implantation









AT LEAST 20% OF KNEE REPLACEMENTS SHOULD BE PARTIALS?

Research shows that surgeons utilizing Partial Knee Arthroplasty (PKA) for at least **20% of their annual knee arthroplasties experienced a significant decrease** in their revision rate.¹⁸

In fact, a study by Badawy, M. et al. found a **lower risk of revision in hospitals performing more than 40 PKAs per year** compared to those performing under 10 PKAs per year.¹⁹

When using criteria published by Kozinn & Scott²⁰ in 1989 only 5% of patients are candidates for PKA.²¹ This may **partly explain why there is low utilization of PKA today, with it only being used for 8% of knee replacements worldwide.**^{1,22,23}

In 2015, Scott revisited the 1989 criteria.²⁴ Using published data, he and five co-authors concluded that some of the **original** contraindications from the 1989 study are no longer considered, thereby increasing patient candidacy.

More recently, in 2009, one study showed that 47.6% of all knee replacement patients are candidates for PKA.²⁵

A randomized, controlled study also showed that significantly **more PKA patients would choose to have their operation again,** compared to TKA patients.²⁶

Other benefits of PKA vs TKA include retention of the ACL, which is reported to result in improved proprioception, ^{27,28} better range of motion, ^{29,30} procedural savings, ²⁵ shorter hospital stays³¹ and a lower risk of postoperative complications. ³²

INTERACTIVE TRAING



Persona Partial Knee Courses

These courses provide the opportunity to learn more about the indications for the Persona Partial Knee and to practice the surgical technique in a hands-on environment with expert faculty.



vLearning

The 'video learning' platform provides an interactive and immersive way to learn more about the Persona Partial Knee. It acts as a 'one-stop-resource' for product & procedure topics.

As you watch the surgical video module, you can opt to select from many relevant resources via sidebar pop-ups or through the quick selection menu.

vLearning is on zbinetwork.com



Surgeon-to-Surgeon (S2S) Visitation

Visitations are available for health care professionals at any level at approved Persona Partial Knee facilities. The visitation includes a discussion about the surgical steps, design rationale, system benefits, and the process for implant and instrument assembly. Surgeons will also be able to better visualize the surgical flow, having spent time with Zimmer Biomet faculty viewing live surgery.



Virtual Reality

Virtual Reality (VR) is one of the newest learning options offered by the Zimmer Biomet Institute.

VR provides you with a variety of learning, training, and practice opportunities.
The immersive nature of VR is capable of delivering a learning experience that is effective and time saving.

Please contact your local sales associate for more information.



References

- * The M/G trademark is owned by Smith & Nephew
- Data on file at Zimmer Biomet. Based on Eucomed IOC RECON Survey 2018.
- 2. Decision Resources Group Large Joint Reconstructve Implants Market Insights, 2020.
- 3. Data on File at Zimmer Biomet. Global Knee Market Model 2020.
- 4. Zimmer ZRR_WA_2409_11.
- 5. Zimmer ZRR_WA_2537_12.
- 6. Zimmer TM1140.98
- 7. Zimmer ZRR_WA_2403_11.
- 8. Zimmer ZRR_WA_2551_12.
- Foran, JR. et al. Long-term Survivorship and Failure Modes of Unicompartmental Knee Arthroplasty. Clin Orthop Relat Res. 2013 Jan; 471(1): 102–108.
- Berger, RA. et al. Results of unicompartmental knee arthroplasty at a minimum of ten years of follow-up. J Bone Joint Surg Am. 2005 May;87(5):999-1006.
- Persona® Partial Knee Clinical Outcomes Study ANNUAL REPORT K.CR.I.G.16.16. Internal data on file at Zimmer Biomet. Sept 2019.
- Bischoff JE, Argenson J-N, Della Valle CJ, Thienpont E. Morphologic Assessment of Tibial Tray Design in Partial Knee Arthroplasty. ORS [submitted]. New Orleans, LA 2018.
- 13. Shepard, M. F., *et al.* Influence of Cement Technique on the Interface Strength of Femoral Components. Orthopaedics and Related Research, December, 2000,381:26-35.
- Keller, John C., et al. Factors affecting surgical alloy/ bone cement interface adhesion. Journal of Biomedical Materials Research, 1980, vol. 14,639-651.
- Kavanaugh A, et al. Factors Influencing the Initial Strength of the Tibial Tray-Cement Interface Bond. Bone Joint J 2013 vol. 95-B no. SUPP 34 98.
- Cawley DT, et al. Cementing Techniques for the Tibial Component in Primary Total Knee Replacement. Bone Joint J 2013;95-B:295-300.

All content herein is protected by copyright, trademarks and other intellectual property rights, as applicable, owned by or licensed to Zimmer Biomet or its affiliates unless otherwise indicated, and must not be redistributed, duplicated or disclosed, in whole or in part, without the express written consent of Zimmer Biomet.

This material is intended for health care professionals, the Zimmer Biomet sales force and Zimmer Biomet employees. Distribution to any other recipient is prohibited.

For product information, including indications, contraindications, warnings, precautions, potential adverse effects and patient counseling information, see the package insert and zimmerbiomet.com.

Check for country product clearances and reference product specific instructions for use. Not for distribution in France.

©2020 Zimmer Biomet

- Miller MA, et al. Loss of Cement-bone Interlock in Retrieved Tibial Components from Total Knee Arthroplasties. Clinical Orthopaedics and Related Research. January 2014, 472:304-313.
- Liddle et al. Optimal usage of unicompartmental knee arthroplasty A STUDY OF 41 986 CASES FROM THE NATIONAL JOINT REGISTRY FOR ENGLAND AND WALES. Bone Joint J 2015;97-B:1506-11.
- Badawy, M et al. Higher revision risk for unicompartmental knee arthroplasty in low-volume hospitals Data from 5,791 cases in the Norwegian Arthroplasty Register. Acta Orthopaedica 2014; 85 (4): 342–347.
- Kozinn, S and Scott, R. Current Concepts Review Unicondylar Knee Arthroplasty. The Journal of Bone and Joint Surgery. VOL. 71-A, NO. I. January 1989.
- Pandit, H, et al. Unnecessary contraindications for mobilebearing unicompartmental knee replacement. J Bone Joint Surg [Br] 2011;93-B:622-8
- 22. European Millennium Report Study 2013.
- 23. US Millennium Study 2014.
- 24. Berend, KR, et al. Consensus Statement on Indications and Contraindications for Medial Unicompartmental Knee Arthroplasty. J Surg Orthop Adv. 2015 Winter;24(4):252-6.
- Willis-Owen CA, et al. Unicondylar knee arthroplasty in the UK National Health Service: An analysis of candidacy, outcome and cost efficacy. Knee. 2009 Dec;16(6):473–8.
- Beard D, Price A, Davies L, et al. A Multicentre Randomised Study Comparing Total or Partial Knee Replacement – One Year Results of The Topkat Trial. BASK. Liverpool, UK 2016.
- Pritchett, JW. Patients Prefer A Bicruciate-Retaining or the Medial Pivot Total Knee Prosthesis. Journal of Arthoplasty. Vol. 26 No. 2 2011.
- 28. Katayama, M. et al. Proprioception and Performance After Anterior Cruciate Ligament Rupture. International Orthopaedics (SICOT)
- 29. Amin A, et al. Unicompartmental or Total Knee Replacement? A Direct Comparative Study of Survivorship and Clinical Outcome at Five Years. JBJS Br. 2006; 88-B; Suppl 1, 100.
- Jones, GG, et al. Gait comparison of unicompartmental and total knee arthroplasties with healthy controls. Bone Joint J 2016;(10 SupplB):16–21.
- Lombardi, A. et al. Is Recovery Faster for Mobile-bearing Unicompartmental than Total Knee Arthroplasty? Clinical Orthopedics and Related Research. 467:1450-57. 2009.
- 32. Brown, NM, et al. Total Knee Arthroplasty Has Higher Postoperative Morbidity Than Unicompartmental Knee Arthroplasty: A Multicenter Analysis. The Journal of Arthroplasty. (2012).

